

Decorrelation estimates for a 1D tight binding model in the localized regime

Trinh Tuan Phong

December 20, 2012

Abstract

In this article, we prove decorrelation estimates for the eigenvalues of a 1D discrete tight binding model near two distinct energies in the localized regime. Consequently, with an arbitrary, fixed number n , the asymptotic independence for local level statistics near n distinct energies is obtained.

1 Introduction

The present paper deals with the following lattice Hamiltonian with off-diagonal disorder in dimension 1: for $u = \{u(n)\}_{n \in \mathbb{Z}} \in l^2(\mathbb{Z})$, set

$$(H_\omega u)(n) = \omega_n(u(n) - u(n+1)) - \omega_{n-1}(u(n-1) - u(n)). \quad (1.1)$$

This model appears in the description of waves (light, acoustic waves, etc) which propagate through a disordered, discrete medium (c.f. [1]).

We can see $\{\omega_n\}_{n \in \mathbb{Z}}$ in this model as weights of bonds of the lattice $\mathbb{Z}$.

Throughout this article, we assume that $\omega := \{\omega_n\}_{n \in \mathbb{Z}}$ are non-negative i.i.d. random variables (r.v.'s for short) with a bounded, compactly supported density ρ .

In addition, from Section 1 to Section 3, we assume more that $\omega_n \in [\alpha_0, \beta_0]$ for all $n \in \mathbb{Z}$ where $\beta_0 > \alpha_0 > 0$. In Section 4, we will comment on relaxing the hypothesis of the lower bound of r.v.'s ω .

It is known that (see [6]):

- the operator H_ω admits an almost sure spectrum $\Sigma := \sigma(H_\omega) = [0, 4\beta_0]$.
- H_ω has an integrated density of states defined as follows:
 ω -a.s., the following limit exists and is ω independent:

$$N(E) := \lim_{|\Lambda| \rightarrow +\infty} \frac{\{\text{e.v. of } H_\omega \text{ less than } E\}}{|\Lambda|} \text{ for a.e. } E. \quad (1.2)$$

As a direct consequence of the Wegner estimate (see Theorem 2.1 in Section 2), $N(E)$ is defined everywhere in $\mathbb{R}$ and absolutely continuous w.r.t. Lebesgue measure with a bounded derivative $\nu(E)$ called the density of states of H_ω .

In the present paper, we follow a usual way to study various statistics related to random operators. We restrict the operator H_ω on some interval $\Lambda \subset \mathbb{Z}$ of finite length with some boundary condition and obtain a finite-volume operator which is denoted by $H_\omega(\Lambda)$. Then, we study diverse statistics for this operator in the limit when $|\Lambda|$ goes to infinity.

Throughout this paper, the boundary condition to define $H_\omega(\Lambda)$ is always the periodic boundary condition. For example, if $\Lambda = [1, N]$, the operator $H_\omega(\Lambda)$ is a self-adjoint $N \times N$ matrix of the following form:

$$\begin{pmatrix} \omega_N + \omega_1 & -\omega_1 & 0 & \dots & 0 & -\omega_N \\ -\omega_1 & \omega_1 + \omega_2 & -\omega_2 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & \omega_{N-2} + \omega_{N-1} & -\omega_{N-1} \\ -\omega_N & 0 & 0 & \dots & -\omega_{N-1} & \omega_{N-1} + \omega_N \end{pmatrix}$$

For $L \in \mathbb{N}$, let $\Lambda = \Lambda_L := [-L, L]$ be a large interval in $\mathbb{Z}$ and $|\Lambda| := (2L+1)$ be its cardinality.

We will denote the eigenvalues of $H_\omega(\Lambda)$ ordered increasingly and repeated according to multiplicity by $E_1(\omega, \Lambda) \leq E_2(\omega, \Lambda) \leq \dots \leq E_{|\Lambda|}(\omega, \Lambda)$.

Let I be the localized region in Σ where the finite-volume fractional-moment criteria for localization are satisfied for the finite-volume operators $H_\omega(\Lambda)$ when $|\Lambda|$ large enough (see Section 2 for more details). In this region, the spectrum of H_ω is pure point and the corresponding eigenfunctions are exponentially decay.

Pick E a positive energy in I with $\nu(E) > 0$, and define the local level statistics near E as follows

$$\Xi(\xi, E, \omega, \Lambda) = \sum_{n=1}^{|\Lambda|} \delta_{\xi_n}(E, \omega, \Lambda)(\xi) \quad (1.3)$$

where

$$\xi_n(E, \omega, \Lambda) = |\Lambda| \nu(E) (E_n(\omega, \Lambda) - E). \quad (1.4)$$

For the model (1.1), It is known that the weak limit of the above point process is a Poisson point process:

Theorem 1.1. *[[6], [8]] Assume that E is a positive energy in I with $\nu(E) > 0$.*

Then, when $|\Lambda| \rightarrow +\infty$, the point process $\Xi(\xi, E, \omega, \Lambda)$ converges weakly to a

Poisson point process with the intensity 1 i.e., for $(U_j)_{1 \leq j \leq J}$, $U_j \subset \mathbb{R}$ bounded measurable and $U_{j'} \cap U_j = \emptyset$ if $j \neq j'$ and $(k_j)_{1 \leq j \leq J} \in \mathbb{N}^J$, we have

$$\lim_{|\Lambda| \rightarrow +\infty} \left| \mathbb{P} \left(\begin{cases} \#\{j; \xi_j(E, \omega, \Lambda) \in U_1\} & = k_1 \\ \vdots & \vdots \\ \#\{j; \xi_j(E, \omega, \Lambda) \in U_J\} & = k_J \end{cases} \right) - \prod_{j=1}^J \frac{|U_j|^{k_j}}{k_j!} e^{-|U_j|} \right| = 0.$$

Recently, for the 1D discrete Anderson model, Klopp [9] showed more that if we pick two fixed, distinct energies E and E' in the localized regime, their two corresponding point processes $\Xi(\xi, E, \omega, \Lambda)$ and $\Xi(\xi, E', \omega, \Lambda)$ converge weakly, respectively to two independent Poisson point processes. In other words, the limits of $\Xi(\xi, E, \omega, \Lambda)$ and $\Xi(\xi, E', \omega, \Lambda)$ are stochastically independent.

From the argument in [9], the above statement holds true if one can prove a so-called decorrelation estimate.

That is exactly what we want to carry out here for the 1D discrete lattice Hamiltonian with off-diagonal disorder (1.1). Our decorrelation estimate is the following:

Theorem 1.2. *Let E, E' be two positive, distinct energies in the localized regime. Pick $\beta \in (1/2, 1)$ and $\alpha \in (0, 1)$. Then, for any $c > 0$, there exists $C > 0$ such that, for L large enough and $cL^\alpha \leq l \leq L^\alpha/c$, one has*

$$\mathbb{P} \left(\left\{ \sigma(H_\omega(\Lambda_l)) \cap (E + L^{-1}(-1, 1)) \neq \emptyset \right\} \right) \leq C(l/L)^2 e^{(\log L)^\beta}.$$

This decorrelation estimate means that, up to a sub-polynomial error, the probability of obtaining simultaneously two eigenvalues near two distinct energies is bounded by the product of the estimates given by Wegner estimate for each of these two energies. Roughly speaking, two eigenvalues of our model near two distinct energies behave like two independent random variables.

Thanks to Theorem 1.2 and an argument in [9], the asymptotic independence of the weak limits of $\Xi(\xi, E, \omega, \Lambda)$ and $\Xi(\xi, E', \omega, \Lambda)$ associated to the model (1.1) is obtained:

Theorem 1.3. *Pick two positive, distinct energies E and E' in the localized regime such that $\nu(E) > 0$ and $\nu(E') > 0$.*

When $|\Lambda| \rightarrow +\infty$, the point processes $\Xi(\xi, E, \omega, \Lambda)$, and $\Xi(\xi, E', \omega, \Lambda)$ converge weakly respectively to two independent Poisson processes on $\mathbb{R}$ with intensity the Lebesgue measure.

That is, for $(U_j)_{1 \leq j \leq J}$, $U_j \subset \mathbb{R}$ bounded measurable and $U_{j'} \cap U_j = \emptyset$ if $j \neq j'$ and $(k_j)_{1 \leq j \leq J} \in \mathbb{N}^J$ and $(U'_j)_{1 \leq j \leq J'}$, $U'_j \subset \mathbb{R}$ bounded measurable and

$U'_{j'} \cap U'_j = \emptyset$ if $j \neq j'$ and $(k'_j)_{1 \leq j \leq J'} \in \mathbb{N}^{J'}$, the quantity

$$\mathbb{P} \left(\begin{pmatrix} \#\{j; \xi_j(E, \omega, \Lambda) \in U_1\} & = k_1 \\ \vdots & \vdots \\ \#\{j; \xi_j(E, \omega, \Lambda) \in U_j\} & = k_J \\ \#\{j; \xi_j(E', \omega, \Lambda) \in U'_1\} & = k'_1 \\ \vdots & \vdots \\ \#\{j; \xi_j(E', \omega, \Lambda) \in U'_J\} & = k'_J \end{pmatrix} \right) - \prod_{j=1}^J \frac{|U_j|^{k_j}}{k_j!} e^{-|U_j|} \prod_{i=1}^{J'} \frac{|U'_i|^{k'_i}}{k'_i!} e^{-|U'_i|}$$

converges to 0 as $|\Lambda|$ goes to infinity.

To prove Theorem 1.2 for the model (1.1), we follow the strategy introduced in [9]. The key point of the proof of decorrelation estimates for the 1D discrete Anderson model in [9] is to derive that the gradient with respect to ω of two eigenvalues $E(\omega)$ and $E'(\omega)$ near two distinct energies E and E' are not co-linear with a good probability. In the discrete Anderson case, this statement will hold true if the gradients of $E(\omega)$ and $E'(\omega)$ are distinct. In deed, the gradients of $E(\omega)$ and $E'(\omega)$ have non-negative components and their l^1 -norm are always equal to 1. So, if they are co-linear, they should be the same.

Unfortunately, the l^1 -norm of the gradient of an eigenvalue of our finite-volume operator $H_\omega(\Lambda)$ is not a constant w.r.t. ω anymore (it is even not bounded from below by a positive constant uniformly w.r.t. ω). Moreover, to prove the above key point for the discrete Anderson model, [9] exploits the diagonal structure of the potential which can not be used for the present case. So, a different approach in the proof is needed to carry out Theorem 1.2. This approach is contained in Lemma 3.3.

In addition, according to [8], a result similar to Theorem 1.3 can be obtained for our model:

Theorem 1.4 (Theorem 1.12, [8]). *Pick $0 < E_0 \in I$ such that the density of states ν is continuous and positive at E_0 .*

Consider two sequences of positive energies, say $(E_\Lambda)_\Lambda$, $(E'_\Lambda)_\Lambda$ such that

$$1. \quad E_\Lambda \xrightarrow{\Lambda \rightarrow \mathbb{Z}^d} E_0 \text{ and } E'_\Lambda \xrightarrow{\Lambda \rightarrow \mathbb{Z}^d} E_0,$$

$$2. \quad |\Lambda| |N(E_\Lambda) - N(E'_\Lambda)| \xrightarrow{\Lambda \rightarrow \mathbb{Z}^d} +\infty.$$

Then, the point processes $\Xi(\xi, E_\Lambda, \omega, \Lambda)$ and $\Xi(\xi, E'_\Lambda, \omega, \Lambda)$ converges weakly respectively to two independent Poisson point processes in $\mathbb{R}$ with intensity the Lebesgue measure.

In Theorem 1.4, instead of fixing two distinct energies E and E' , one considers two sequences of energies $\{E_\Lambda\}$, $\{E'_\Lambda\}$ which tend to each other as $|\Lambda| \rightarrow \infty$. In addition, we assume that, roughly speaking, there are sufficiently many

eigenvalues of $H_\omega(\Lambda)$ between E_Λ and E'_Λ as Λ large. Then, we can derive again the asymptotic independence of two point processes associated to E_Λ and E'_Λ .

Besides, it is known that the existence of an integrated density of states defined as in (1.2) implies that the average distance (mean spacing) between eigenlevels is of order $|\Lambda|^{-1}$.

Thus, Theorem 1.4 shows that, in the localized regime, eigenvalues separated by a distance that is asymptotically infinite with respect to the mean spacing between eigenlevels behave like independent random variables. In other words, there are no interactions between distinct eigenvalues, except at a very short distance.

Finally, in Section 5, we will see that a similar statement to Theorem 1.3 holds true as we consider not only two but any fixed number of distinct energies.

Notation: In the present paper, we use Dirac's notations: $|x\rangle = \delta_x$, and $|x\rangle\langle x| = \langle \delta_x, \cdot \rangle \delta_x$ is the projection operator on δ_x .

2 Preliminaries

We recall here a Wegner-type estimate and a Minami-type estimate for the model (1.1) which are essentially important for the proofs of Poisson statistics as well as those of decorrelation estimates.

Theorem 2.1. [*Wegner estimate, Theorem 2.1, [6]*]

$$\mathbb{P}(\text{dist}(E, \sigma(H_\omega(\Lambda))) \leq \epsilon) \leq \frac{2d \|s\rho(s)\|_\infty}{E - \epsilon} \epsilon |\Lambda|$$

for all intervals $\Lambda \subset \mathbb{Z}$ and $0 < \epsilon < E$.

Roughly speaking, the above Wegner estimate means that the probability that we can find an eigenvalue of the finite-volume operator $H_\omega(\Lambda)$ in a interval (suppose that the length of this interval is small) is bounded by the length of this interval times $|\Lambda|$.

Following is the Minami estimate:

Theorem 2.2. [*Minami estimate, Theorem 3.1, [6]*] *There exists $C > 0$ such that, for all intervals $J = [a, b] \subset (0, +\infty)$, and $\Lambda \subset \mathbb{Z}$, we have*

$$\mathbb{P}(\#\{\sigma(H_\omega(\Lambda)) \cap J\} \geq 2) \leq C |J|^2 |\Lambda|^2.$$

It is not hard to realize a similarity between the Minami estimate and the decorrelation estimate in Theorem 1.2: the Minami estimate can be interpreted as a decorrelation estimate for close together eigenvalues and can be used to get the counterparts of Theorem 1.1 and 1.3 when E and E' are closer and closer to each other as $|\Lambda| \rightarrow \infty$ (c.f. [8]).

We should remark that although the almost sure spectrum Σ of the operator (1.1) contains the point 0 (the bottom of Σ), the above Wegner and Minami estimates do not work at 0. That is why, in the present paper, we will only consider energies who are strictly positive in Σ .

We would like to remind readers of the precise definition of the localized regime or the region of localization.

Proposition 2.3. (c.f. [2], [9]) *Let I be the region of Σ where the finite volume fractional moment criteria of [2] for $H_\omega(\Lambda)$ are verified for Λ sufficiently large.*

Then, there exists $\nu > 0$ such that, for any $p > 0$, there exists $q > 0$ and $L_0 > 0$ such that, for $L \geq L_0$, with probability larger than $1 - L^{-p}$, if

1. $\varphi_{n,\omega}$ is a normalized eigenvector of $H_\omega(\Lambda_L)$ associated to an energy $E_{n,\omega} \in I$,
2. $x_{n,\omega} \in \Lambda_L$ is a maximum of $x \mapsto |\varphi_{n,\omega}(x)|$ in Λ_L ,

then, for $x \in \Lambda_L$, one has

$$|\varphi_{n,\omega}(x)| \leq L^q e^{-\nu|x-x_{n,\omega}|}.$$

The point $x_{n,\omega}$ is called a localization center for $\varphi_{n,\omega}$ or $E_{n,\omega}$.

Remark that an eigenvalue of $H_\omega(\Lambda)$ can have more than one localization center. Nevertheless, it is not hard to check that, under assumptions of Proposition 2.3, all localization centers associated to one eigenvalue are contained in a disk of radius $C \log L$ with some constant C .

Hence, to define the localization center uniquely for an eigenvalue, we can order these centers lexicographically and choose the localization center associated to an eigenvalue to be the largest one.

3 Proof of the decorrelation estimate

Pick two distinct, positive energies E, E' in I . Let $J_L = E + L^{-1}[-1, 1]$ and $J'_L = E' + L^{-1}[-1, 1]$ with L large.

Lemma 3.1. *Suppose that $\omega \mapsto E(\omega)$ is the only eigenvalue of $H_\omega(\Lambda)$ in J_L . Then*

1. $E(\omega)$ is simple and $\omega \mapsto E(\omega)$ is real analytic, and if $\omega \mapsto \varphi(\omega)$ is the corresponding real-valued, normalized eigenvector, it is real analytic in ω .
2. $\|\nabla_\omega E(\omega)\|_{l^1} = 2 \sum_{\gamma \in \Lambda} \|\Pi_\gamma \varphi\|^2$ where $\Pi_\gamma = \frac{1}{2}|\gamma - (\gamma + 1)\rangle\langle\gamma - (\gamma + 1)|$ is a projection in $l^2(\Lambda)$.

3. $\text{Hess}_\omega E(\omega) = (h_{\gamma,\beta})_{\gamma,\beta}$ where

- $h_{\gamma,\beta} := -4\text{Re}\langle (H_\omega(\Lambda) - E(\omega))^{-1}\psi_\gamma, \psi_\beta \rangle$.
- $\psi_\gamma := \langle \Pi_\gamma \varphi, \varphi \rangle \varphi - \Pi_\gamma \varphi = -\Pi_{\langle \varphi \rangle^\perp}(\Pi_\gamma \varphi)$ where $\Pi_{\langle \varphi \rangle^\perp}$ is the orthogonal projection on $\langle \varphi \rangle^\perp$.

Proof. (1) is true from the standard perturbation theory (see [10]).

Now we will prove (2). Starting from the eigenequation:

$$H_\omega(\Lambda)\varphi = E(\omega)\varphi \quad (3.1)$$

we have, for all $\gamma \in \Lambda$,

$$\begin{aligned} \partial_{\omega_\gamma} E(\omega) &= \langle \partial_{\omega_\gamma}(H_\omega(\Lambda)\varphi), \varphi \rangle + \langle H_\omega(\Lambda)\varphi, \partial_{\omega_\gamma}\varphi \rangle \\ &= \langle \partial_{\omega_\gamma}(H_\omega(\Lambda))\varphi, \varphi \rangle + \langle H_\omega(\Lambda)\partial_{\omega_\gamma}\varphi, \varphi \rangle + \langle H_\omega(\Lambda)\varphi, \partial_{\omega_\gamma}\varphi \rangle \\ &= \langle \partial_{\omega_\gamma}(H_\omega(\Lambda))\varphi, \varphi \rangle + \langle \partial_{\omega_\gamma}\varphi, H_\omega(\Lambda)\varphi \rangle + \langle H_\omega(\Lambda)\varphi, \partial_{\omega_\gamma}\varphi \rangle \\ &= \langle \partial_{\omega_\gamma}(H_\omega(\Lambda))\varphi, \varphi \rangle + E(\omega)(\langle \partial_{\omega_\gamma}\varphi, \varphi \rangle + \langle \varphi, \partial_{\omega_\gamma}\varphi \rangle) \end{aligned}$$

where the last two equalities come from the symmetry of $H_\omega(\Lambda)$ and (3.1).

Noting that

$$\langle \partial_{\omega_\gamma}\varphi, \varphi \rangle + \langle \varphi, \partial_{\omega_\gamma}\varphi \rangle = 2\partial_{\omega_\gamma}\|\varphi\|^2 = 0.$$

Hence,

$$\partial_{\omega_\gamma} E(\omega) = \langle \partial_{\omega_\gamma}(H_\omega(\Lambda))\varphi, \varphi \rangle = 2\|\Pi_\gamma \varphi\|^2 \geq 0. \quad (3.2)$$

Besides, it is easy to check that $\Pi_\gamma = \Pi_\gamma^* = \Pi_\gamma^2$. Hence, Π_γ is an orthogonal projection.

From (3.2), we have the following important equality:

$$\sum_{\gamma \in \Lambda} \omega_\gamma \partial_{\omega_\gamma} E(\omega) = E(\omega) \quad (3.3)$$

which characterize the form of our operator.

Finally, we give a proof for (3). By differentiating both sides of (3.2) with respect to ω_γ , we have

$$\begin{aligned} \partial_{\omega_\gamma}^2 E(\omega) &= 2\langle \partial_{\omega_\gamma}(\Pi_\gamma \varphi), \varphi \rangle + 2\langle \Pi_\gamma \varphi, \partial_{\omega_\gamma}\varphi \rangle \\ &= 2\langle \Pi_\gamma \partial_{\omega_\gamma}\varphi, \varphi \rangle + 2\langle \Pi_\gamma \varphi, \partial_{\omega_\gamma}\varphi \rangle = 4\text{Re}\langle \Pi_\gamma \varphi, \partial_{\omega_\gamma}\varphi \rangle \end{aligned} \quad (3.4)$$

Next, we will compute $\partial_{\omega_\gamma}\varphi$.

Differentiating both sides of (3.1) with respect to ω_γ to get

$$\begin{aligned} (\partial_{\omega_\gamma} H_\omega(\Lambda))\varphi + H_\omega(\Lambda)\partial_{\omega_\gamma}\varphi &= \partial_{\omega_\gamma} E(\omega)\varphi + E(\omega)\partial_{\omega_\gamma}\varphi \\ &= 2\langle \Pi_\gamma \varphi, \varphi \rangle \varphi + E(\omega)\partial_{\omega_\gamma}\varphi \end{aligned}$$

Therefore,

$$[H_\omega(\Lambda) - E(\omega)]\partial_{\omega_\gamma}\varphi = 2\langle \Pi_\gamma \varphi, \varphi \rangle \varphi - (\partial_{\omega_\gamma} H_\omega(\Lambda))\varphi = 2\left(\langle \Pi_\gamma \varphi, \varphi \rangle \varphi - \Pi_\gamma \varphi\right).$$

Observe that $\psi_\gamma := \langle \Pi_\gamma \varphi, \varphi \rangle \varphi - \Pi_\gamma \varphi \in \langle \varphi \rangle^\perp$, and $[H_\omega(\Lambda) - E(\omega)]$ is invertible in the subspace $\langle \varphi \rangle^\perp$ of $l^2(\Lambda)$, we get

$$\partial_{\omega_\gamma} \varphi = 2(H_\omega(\Lambda) - E(\omega))^{-1} (\langle \Pi_\gamma \varphi, \varphi \rangle \varphi - \Pi_\gamma \varphi). \quad (3.5)$$

From (3.4) and (3.5), we obtain

$$\partial_{\omega_\gamma}^2 E(\omega) = 4\text{Re} \langle \Pi_\gamma \varphi, (H_\omega(\Lambda) - E(\omega))^{-1} (\langle \Pi_\gamma \varphi, \varphi \rangle \varphi - \Pi_\gamma \varphi) \rangle.$$

Thanks to (3.5), we have $2(H_\omega(\Lambda) - E(\omega))^{-1} (\langle \Pi_\gamma \varphi, \varphi \rangle \varphi - \Pi_\gamma \varphi)$ is orthogonal to φ . We therefore infer that

$$\partial_{\omega_\gamma}^2 E(\omega) = -4\text{Re} \langle (H_\omega(\Lambda) - E(\omega))^{-1} \psi_\gamma, \psi_\gamma \rangle.$$

Repeating this argument, it is not hard to prove that

$$\partial_{\omega_\gamma \omega_\beta}^2 E(\omega) = -4\text{Re} \langle (H_\omega(\Lambda) - E(\omega))^{-1} \psi_\gamma, \psi_\beta \rangle$$

for all γ, β . So, we have Lemma 3.1 proved. $\square$

Next, we would like to state here a "lower bound" for normalized eigenfunctions of $H_\omega(\Lambda)$ in the sense that there exists a large subset J in Λ such that the components $(u(k))_{k \in J}$ of a normalized eigenfunction u of $H_\omega(\Lambda)$ can not be too small.

Lemma 3.2. *Let $\Lambda = [-L, L]$ be a large cube in $\mathbb{Z}$ and $\beta \in (1/2, 1)$. Then, there exists a point k_0 in Λ and a positive constant c such that $u^2(k) + u^2(k+1) \geq e^{-L^\beta}$ for all $|k - k_0| \leq cL^\beta$ when L is large enough.*

Proof. We rewrite the eigenequation corresponding to the eigenvector u and eigenvalue E at the point n by means of the transfer matrix

$$\begin{pmatrix} u(n+1) \\ u(n) \end{pmatrix} = \begin{pmatrix} \frac{\omega_n + \omega_{n-1} - E}{\omega_n} & \frac{-\omega_{n-1}}{\omega_n} \\ 1 & 0 \end{pmatrix} \begin{pmatrix} u(n) \\ u(n-1) \end{pmatrix}.$$

Now, we denote the transfer matrix

$$\begin{pmatrix} \frac{\omega_n + \omega_{n-1} - E}{\omega_n} & \frac{-\omega_{n-1}}{\omega_n} \\ 1 & 0 \end{pmatrix}$$

by $T(n, E)$ and denote by $v(n)$ the column vector $= (u(n+1), u(n))^t$. Hence, for all n greater than m we have $v(n) = T(n, E) \cdots T(n-m+1, E)v(m)$. Remark that the transfer matrices are invertible. Moreover, they and their inverse matrices are uniformly bounded by a constant C depending only on the support of ω_j , because $\omega_j \in [\alpha_0, \beta_0]$ with $0 < \alpha_0 < \beta_0$. Thus,

$$|v(n)| \leq C^{|n-m|} |v(m)| = e^{\log C |n-m|} |v(m)| = e^{\eta |n-m|} |v(m)|$$

for all n, m in Λ .

Assume that $|v(k_0)|$ is the maximum of $|v(n)|$. Hence,

$$|v(k_0)| \geq \frac{1}{\sqrt{2L}}$$

from the fact that $\sum_{j \in \Lambda} |v(j)|^2 = 2$. Thus, for any $\kappa > 0$, we obtain

$$|v(k)| \geq \frac{1}{\sqrt{2L}} e^{-\eta|k-k_0|} \geq e^{-\frac{\kappa\eta L^\beta}{2}}$$

when $|k - k_0| \leq \kappa L^\beta$ and L sufficiently large.

Hence, we have

$$u^2(k) + u^2(k+1) \geq e^{-\kappa\eta L^\beta}$$

for all $|k - k_0| \leq \kappa L^\beta$. So, by choosing $\kappa = \frac{1}{\eta}$, we have Lemma 3.2 proved. $\square$

The following lemma is the main ingredient of the proof of the decorrelation estimate as well as the heart of the present paper:

Lemma 3.3. *Let E, E' be two positive energies in the localized regime and $\beta > 1/2$. Let $c_1, c_2 > 0$ and denote by $\mathbb{P}^*$ the probability of the following event (called $(*)$):*

There exists two simple eigenvalues of $H_\omega(\Lambda_L)$, say $E(\omega), E'(\omega)$ such that $|E(\omega) - E| + |E'(\omega) - E'| \leq e^{-L^\beta}$ and

$$\|\nabla_\omega(c_1^2 E(\omega) - c_2^2 E'(\omega))\|_1 \leq c_1^2 e^{-L^\beta}.$$

Then, there exists $c > 0$ such that

$$\mathbb{P}^* \leq e^{-cL^{2\beta}}$$

Remark: There is a slightly but really important difference between the above lemma and Lemma 2.4 in [9] where $c_1 = c_2 = 1$.

In fact, in the proof of the decorrelation estimate for our model, we will use the above lemma with c_1^2, c_2^2 are respectively $\frac{1}{E}, \frac{1}{E'}$ which are two distinct, positive numbers. This difference results from the lack of the normalization of $\|\nabla E(\omega)\|_1$ for our model.

Moreover, we will see in the remark at the end of this section that, for the model (1.1), if $c_1 = c_2$, P^* is exactly equal to 0 when L large enough.

We will skip for a moment the proof of Lemma 3.3 and recall how to use the above lemma to complete the proof of Theorem 1.2. This part can be found in [9]. We repeat it here (with tiny modifications) just for the readers convenience.

Proof. Recall that $J_L = E + L^{-1}[-1, 1]$ and $J'_L = E' + L^{-1}[-1, 1]$ with L large.

Let $cL^\alpha \leq l \leq L^\alpha/c$ where c is fixed. By the Minami estimate, we have

$$\mathbb{P}(\#\{\sigma(H_\omega(\Lambda_l)) \cap J_L\} \geq 2 \text{ or } \#\{\sigma(H_\omega(\Lambda_l)) \cap J'_L\} \geq 2) \leq (l/L)^2.$$

Hence, it is sufficient to prove that $\mathbb{P}_0 \leq C(l/L)^2 e^{(\log L)^\beta}$ where

$$\mathbb{P}_0 := \mathbb{P}(\#\{\sigma(H_\omega(\Lambda_l)) \cap J_L\} = 1; \#\{\sigma(H_\omega(\Lambda_l)) \cap J'_L\} = 1). \quad (3.6)$$

The crucial idea of proving decorrelation estimates in [9] is to reduce the proof of (3.6) to the proof of a similar estimate where Λ_l is replaced by a much smaller cube, a cube of side length of order $\log L$. Precisely, one has (c.f. Lemma 2.1 and Lemma 2.2 in [9]):

$$\mathbb{P}_0 \leq C(l/L)^2 + C(l/\tilde{l})\mathbb{P}_1$$

where $\tilde{l} = C \log L$ and

$$\mathbb{P}_1 := \mathbb{P}(\#\{\sigma(H_\omega(\Lambda_{\tilde{l}})) \cap \tilde{J}_L\} \geq 1 \text{ and } \#\{\sigma(H_\omega(\Lambda_{\tilde{l}})) \cap \tilde{J}'_L\} \geq 1)$$

where $\tilde{J}_L = E + L^{-1}(-2, 2)$ and $\tilde{J}'_L = E' + L^{-1}(-2, 2)$.

To complete the prove of Theorem 1.2, one proves that

$$\mathbb{P}_1 \leq C(\tilde{l}/L)^2 e^{\tilde{l}^\beta}. \quad (3.7)$$

Combining the Minami estimate and the following lemma

Lemma 3.4. [Lemma 2.3, [9]] *There exists $C > 0$ such that*

$$\|Hess_\omega(E(\omega))\|_{l^\infty \rightarrow l^1} \leq \frac{C}{\text{dist}(E(\omega), \sigma(H_\omega(\Lambda)) \setminus E(\omega))}$$

we obtain that

$$\mathbb{P}\left(\left\{\begin{array}{l} \sigma(H_\omega(\Lambda_{\tilde{l}})) \cap \tilde{J}_L = \{E(\omega)\} \\ \|\text{Hess}_\omega(E(\omega))\|_{l^\infty \rightarrow l^1} \geq \epsilon^{-1} \end{array}\right\}\right) \leq C\epsilon \tilde{l}^2 L^{-1}.$$

Hence, for $\epsilon \in (4L^{-1}, 1)$, one has

$$\mathbb{P}_1 \leq C\epsilon \tilde{l}^2 L^{-1} + \mathbb{P}_\epsilon \quad (3.8)$$

where $\mathbb{P}_\epsilon = \mathbb{P}(\Omega_0(\epsilon))$ and

$$\Omega_0(\epsilon) = \left\{ \begin{array}{l} \sigma(H_\omega(\Lambda_{\tilde{l}})) \cap \tilde{J}_L = \{E(\omega)\} \\ \{E(\omega)\} = \sigma(H_\omega(\Lambda_{\tilde{l}})) \cap (E - C\epsilon, E + C\epsilon) \\ \sigma(H_\omega(\Lambda_{\tilde{l}})) \cap \tilde{J}'_L = \{E'(\omega)\} \\ \{E'(\omega)\} = \sigma(H_\omega(\Lambda_{\tilde{l}})) \cap (E' - C\epsilon, E' + C\epsilon) \end{array} \right\}.$$

On the one hand, keeping the same notations as in [9], one puts $\lambda := \frac{1}{2\alpha\beta^2}e^{-\tilde{l}\beta}$. For γ and γ' in Λ , one defines

$$\Omega_{0,\beta}^{\gamma,\gamma'}(\epsilon) = \Omega_0(\epsilon) \cap \{\omega \mid |J_{\gamma,\gamma'}(E(\omega), E'(\omega))| \geq \lambda\}$$

where $J_{\gamma,\gamma'}(E(\omega), E'(\omega))$ is the Jacobian of the mapping

$$(\omega_\gamma, \omega_{\gamma'}) \mapsto (E(\omega), E'(\omega)).$$

Hence, we can dominate $\mathbb{P}_\epsilon$ by

$$\mathbb{P}_\epsilon \leq \sum_{\gamma \neq \gamma'} \mathbb{P}(\Omega_{0,\beta}^{\gamma,\gamma'}(\epsilon)) + \mathbb{P}_r$$

where $\mathbb{P}_r$ is the probability of the following event

$$\mathcal{D} := \{\omega \mid |J_{\gamma,\gamma'}(E(\omega), E'(\omega))| \leq \lambda \text{ for all } \gamma, \gamma' \in \Lambda_{\tilde{l}}\}.$$

Moreover, from Lemma 2.6 in [9], we know that

$$\mathbb{P}(\Omega_{0,\beta}^{\gamma,\gamma'}(\epsilon)) \leq CL^{-2}\lambda^{-4}.$$

Hence,

$$\mathbb{P}_\epsilon \leq C\tilde{l}^4 L^{-2} \lambda^{-4} + \mathbb{P}_r. \quad (3.9)$$

Combining (3.8), (3.9) and choose $\epsilon := L^{-1}\lambda^{-3}$, we infer that

$$\mathbb{P}_1 \leq C(\tilde{l}/L)^2 e^{\tilde{l}\beta} + \mathbb{P}_r. \quad (3.10)$$

Finally, we will use Lemma 3.3 to estimate $\mathbb{P}_r$.

To do so, for each $\omega \in \mathcal{D}$, we rewrite the Jacobian $J_{\gamma,\gamma'}(E(\omega), E'(\omega))$ as follows:

$$\begin{aligned} J_{\gamma,\gamma'}(E(\omega), E'(\omega)) &= \begin{vmatrix} \partial_{\omega_\gamma} E(\omega) & \partial_{\omega_{\gamma'}} E(\omega) \\ \partial_{\omega_\gamma} E'(\omega) & \partial_{\omega_{\gamma'}} E'(\omega) \end{vmatrix} \\ &= \frac{E(\omega)E'(\omega)}{\omega_\gamma \omega_{\gamma'}} \begin{vmatrix} \frac{1}{E(\omega)} \omega_\gamma \partial_{\omega_\gamma} E(\omega) & \frac{1}{E(\omega)} \omega_{\gamma'} \partial_{\omega_{\gamma'}} E(\omega) \\ \frac{1}{E'(\omega)} \omega_\gamma \partial_{\omega_\gamma} E'(\omega) & \frac{1}{E'(\omega)} \omega_{\gamma'} \partial_{\omega_{\gamma'}} E'(\omega) \end{vmatrix}. \end{aligned}$$

if $E(\omega)$ and $E'(\omega)$ are non-zero.

Observe that the sum over γ or γ' of each entry in the second line of the above 2×2 matrix is equal to 1 by (3.3), we can use the following lemma:

Lemma 3.5 (Lemma 2.5, [9]). *Pick $u, v \in (\mathbb{R}^+)^{2n}$ such that $\|u\|_1 = \|v\|_1 = 1$. Then,*

$$\max_{j \neq k} \left| \begin{vmatrix} u_j & u_k \\ v_j & v_k \end{vmatrix} \right|^2 \geq \frac{1}{4n^5} \|u - v\|_1^2$$

to deduce

$$\|\nabla_\omega(\frac{1}{E}E(\omega) - \frac{1}{E'}E'(\omega))\|_1 \leq e^{-\tilde{l}\beta}$$

provided that

$$|E(\omega) - E| + |E'(\omega) - E'| \leq e^{-\tilde{l}\beta}$$

and $\tilde{l} = O(\log L)$ is sufficiently large.

Thus, thanks to Lemma 3.3, for L sufficiently large, and with the remark that $\tilde{l} \geq c \log L$, $\beta > 1/2$, we have

$$\mathbb{P}_r \leq L^{-2}$$

which completes the proof of Theorem 1.2. $\square$

To complete the present section, we state here the proof of Lemma 3.3:

Proof of Lemma 3.3. Let $\omega \mapsto u(\omega)$ and $\omega \mapsto v(\omega)$ be normalized eigenvectors associated to $E(\omega)$ and $E'(\omega)$. By Lemma 3.1, we have

$$\nabla_\omega E(\omega) = (2\|\Pi_\gamma u\|^2)_{\gamma \in \Lambda} \text{ and } \nabla_\omega E'(\omega) = (2\|\Pi_\gamma v\|^2)_{\gamma \in \Lambda}$$

Now, We introduce the linear operator T from $l^2(\Lambda)$ to $l^2(\Lambda)$ defined as follows

$$Tu(n) = u(n) - u(n+1)$$

where $u = (u(n))_n \in l^2(\Lambda)$. Recall that $\Lambda = \Lambda_L = \mathbb{Z}/L\mathbb{Z}$, i.e. we use periodic boundary conditions here.

Assume that

$$\begin{aligned} c_1^2 e^{-L^\beta} &\geq \|\nabla_\omega(c_1^2 E(\omega) - c_2^2 E'(\omega))\|_1 = \sum_n |(c_1 Tu(n))^2 - (c_2 Tv(n))^2| \\ &= \sum_n |c_1 Tu(n) - c_2 Tv(n)| |c_1 Tu(n) + c_2 Tv(n)|. \end{aligned}$$

Hence, $e^{-L^\beta} \geq \sum_n |Tu(n) - cTv(n)| |Tu(n) + cTv(n)|$ with $c = c_2/c_1 > 0$.

Then, there exists a partition of $\Lambda = \{-L, \dots, L\}$, say $\mathcal{P} \subset \Lambda$ and $\mathcal{Q} \subset \Lambda$ such that $\mathcal{P} \cup \mathcal{Q} = \Lambda$, $\mathcal{P} \cap \mathcal{Q} = \emptyset$ and

- for $n \in \mathcal{P}$, $|Tu(n) - cTv(n)| \leq e^{-L^\beta/2}$
- for $n \in \mathcal{Q}$, $|Tu(n) + cTv(n)| \leq e^{-L^\beta/2}$

In the rest of this proof, we put $v(n) := cv(n)$. This abuse of notation changes nothing thanks to the linearity of the operator T .

From Lemma 3.2, there exists a $c_1 > 0$ and an interval J of the length $c_1 L^\beta$ which satisfies

$$|u(k)|^2 + |u(k+1)|^2 \geq e^{-L^\beta/4} \quad (3.11)$$

for all $k \in J$.

Now, we want to study the properties of the above partition in the interval J .

We assume that

$$\mathcal{P} \cap J = \cup \mathcal{P}_j \text{ and } \mathcal{Q} \cap J = \cup \mathcal{Q}_j$$

where $\mathcal{P}_j$ and $\mathcal{Q}_j$ are intervals in $\mathbb{Z}$.

Step 1: We'll show that the length of any $\mathcal{P}_j$ or $\mathcal{Q}_j$ is inferior or equal to 3. Indeed, assume by the contradiction that there exists an interval $\mathcal{P}_j = \{n-2, n-1, n, n+1, n+2 \dots m\}$ with $|\mathcal{P}_j| = |m-n+2| \geq 4$. So, $m \geq n+2$. First of all, the following relations are true for the points in $\mathcal{P}_j$

$$\begin{aligned} Tu(n-2) &= Tv(n-2) + O(e^{-L^\beta/2}). \\ Tu(n-1) &= Tv(n-1) + O(e^{-L^\beta/2}). \\ Tu(n) &= Tv(n) + O(e^{-L^\beta/2}). \\ Tu(n+1) &= Tv(n+1) + O(e^{-L^\beta/2}). \\ Tu(n+2) &= Tv(n+2) + O(e^{-L^\beta/2}). \end{aligned} \tag{3.12}$$

Combining the eigenequations for u and v at the point $n-1$

$$Eu(n-1) = \omega_{n-1}Tu(n-1) - \omega_{n-2}Tu(n-2) + O(e^{-L^\beta/2}) \tag{3.13}$$

$$E'v(n-1) = \omega_{n-1}Tv(n-1) - \omega_{n-2}Tv(n-2) + O(e^{-L^\beta/2})$$

and relations in 3.12, we obtain

$$Eu(n-1) = E'v(n-1) + O(e^{-L^\beta/2}). \tag{3.14}$$

Similarly, we have

$$Eu(n) = E'v(n) + O(e^{-L^\beta/2}). \tag{3.15}$$

Combining (3.14), (3.15) and the second relation in (3.12), we obtain

$$\left(1 - \frac{E}{E'}\right)u(n) = \left(1 - \frac{E}{E'}\right)u(n-1) + O(e^{-L^\beta/2})$$

which implies that

$$Tu(n-1) \leq Ce^{-L^\beta/2} \tag{3.16}$$

where C is a positive constant depending only on E, E' and the support of ω .

In other words, we showed that, if we have three consecutive points in some interval $\mathcal{P}_j$ ($n-2, n-1, n$ in the above case), then the middle point of those three points always satisfies the inequality (3.16).

Repeating the above argument for other points in $\mathcal{P}_j$, we obtain that

$$Tu(n) \leq Ce^{-L^\beta/2}, \tag{3.17}$$

and

$$Tu(n+1) \leq Ce^{-L^\beta/2}. \quad (3.18)$$

Therefore, combining the equations (3.16), (3.17), (3.18) and the eigenequations for u at the point n and $n+1$

$$\begin{aligned} Eu(n) &= \omega_n Tu(n) - \omega_{n-1} Tu(n-1) + O(e^{-L^\beta/2}), \\ Eu(n+1) &= \omega_{n+1} Tu(n+1) - \omega_n Tu(n) + O(e^{-L^\beta/2}), \end{aligned}$$

we infer that there exists a positive constant C being independent of L such that

$$|u(n)|^2 + |u(n+1)|^2 \leq Ce^{-L^\beta}$$

which contradicts (3.11) if we choose L large enough.

Hence, an interval $\mathcal{P}_j$ or $\mathcal{Q}_j$ can not contain more than four points in $\mathbb{Z}$ and we have the statement in **Step 1** proved.

Step 2: Consider four arbitrarily consecutive points in J , say $n-2, n-1, n, n+1$. The following are all possible configurations for these points with the remark of the equivalent role of $\mathcal{P}_j$ and $\mathcal{Q}_j$.

First case: We consider the case when three of these points belong to an interval $\mathcal{P}_j$ and the other belongs to $\mathcal{Q}_j$, say $n-2, n-1, n \in \mathcal{P}_j$ and $n+1 \in \mathcal{Q}_j$.

First of all, we have the same eigenequations as in (3.12) with respect to $n-2, n-1, n$ and the following

$$Tu(n+1) = -Tv(n+1) + O(e^{-L^\beta/2}).$$

Secondly, we have two eigenequations for u and v at the point $n+1$

$$Eu(n+1) = \omega_{n+1} Tu(n+1) - \omega_n Tu(n) + O(e^{-L^\beta/2}) \quad (3.19)$$

and

$$E'v(n+1) = \omega_{n+1} Tv(n+1) - \omega_n Tv(n) + O(e^{-L^\beta/2})$$

which can be rewritten as follows

$$E'v(n+1) = -\omega_{n+1} Tu(n+1) - \omega_n Tu(n) + O(e^{-L^\beta/2}). \quad (3.20)$$

Thirdly, repeating the argument of **Step 1**, we can obtain (3.14) and (3.15) again. So, combining with two eigenequations for u at the point $n-1$ and n , we will have the following system of linear equations (10 equations, 10 variables):

$$A_0 U = b_0$$

where $U = (u(n-2), u(n-1), \dots, u(n+2), v(n-2), v(n-1), \dots, v(n+2))$, $b_0 = (b_0^j)_{1 \leq j \leq 10}$ with $\|b_0\| \leq ce^{-L^\beta/2}$ and A_0 is the following 10×10 matrix:

$$\begin{pmatrix}
1 & -1 & 0 & 0 & 0 \\
0 & 1 & -1 & 0 & 0 \\
0 & 0 & 1 & -1 & 0 \\
0 & 0 & 0 & 1 & -1 \\
0 & -\omega_{n-1} & \omega_{n-1} + \omega_n - E & -\omega_n & 0 \\
0 & 0 & \frac{E}{E'} & 0 & 0 \\
0 & 0 & -\omega_n & \omega_n + \omega_{n+1} - E & -\omega_{n+1} \\
0 & 0 & \omega_n & \omega_{n+1} - \omega_n & -\omega_{n+1} \\
-\omega_{n-2} & \omega_{n-2} + \omega_{n-1} - E & -\omega_{n-1} & 0 & 0 \\
0 & \frac{E}{E'} & 0 & 0 & 0
\end{pmatrix}
\begin{pmatrix}
1 & -1 & 0 & 0 & 0 \\
0 & 1 & -1 & 0 & 0 \\
0 & 0 & 1 & -1 & 0 \\
0 & 0 & 0 & -1 & 1 \\
0 & 0 & 0 & 0 & 0 \\
0 & 0 & -1 & 0 & 0 \\
0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & E' & 0 \\
0 & 0 & 0 & 0 & 0 \\
0 & -1 & 0 & 0 & 0
\end{pmatrix}$$

Let $\widetilde{A}_0$ be the comatrix of A_0 and ρ_0 be $\max\{1, \|\widetilde{A}_0\|\}$.

We will prove that $|\det A_0| \leq \rho_0 e^{-L^\beta/4}$.

Indeed, assume by the contradiction that $|\det A_0| > \rho_0 e^{-L^\beta/4}$.

Therefore, A_0 has the inverse satisfying

$$\|A_0^{-1}\| \leq \frac{\rho_0}{|\det A_0|} \leq e^{L^\beta/4}.$$

Hence,

$$\|U\| = \|A_0^{-1}b_0\| \leq \|A_0^{-1}\| \|b_0\| \leq ce^{L^\beta/4} e^{-L^\beta/2} \leq ce^{-L^\beta/4}$$

But this is impossible according to (3.11). Consequently, $|\det A_0| \leq \rho_0 e^{-L^\beta/4}$.

Remark that ρ_0 only depends on the support of ω_j and E, E' .

Now, we would like to compute explicitly the determinant of the matrix A_0 which is a polynomial of at most four variables ω_j with $j = n-2, n+1$. Then, combining with the fact that this determinant is small, we infer some restrictive conditions on ω .

Put $A_0 = (a_{ij})$, we'll give here some details of computing the determinant of the matrix A_0 by hand:

(A mathematical software like Maple or Mathematica might be useful for checking the final result of this computation).

First, expand this determinant by its sixth and last column and then by its first column to get

$$|\det A_0| = \omega_{n-2} |\det B_0|$$

where B_0 is the 7×7 matrix defined by

$$\begin{pmatrix} 1 & -1 & 0 & 0 & 1 & -1 & 0 \\ 0 & 1 & -1 & 0 & 0 & 1 & -1 \\ -\omega_{n-1} & \omega_{n-1} + \omega_n - E & -\omega_n & 0 & 0 & 0 & 0 \\ 0 & \frac{E}{E'} & 0 & 0 & 0 & -1 & 0 \\ 0 & -\omega_n & \omega_n + \omega_{n+1} - E & -\omega_{n+1} & 0 & 0 & 0 \\ 0 & \omega_n & \omega_{n+1} - \omega_n & -\omega_{n+1} & 0 & 0 & E' \\ -\frac{E}{E'} & 0 & 0 & 0 & -1 & 0 & 0 \end{pmatrix}$$

Now, we compute the determinant of B_0 .

Take the sixth row minus the fifth row and take the first row plus the last row. Next, multiply the second row by E' and take it plus the sixth row. Finally, expand the determinant of B_0 by the forth, the fifth and the last column to get

$$|\det A_0| = \frac{1}{E'} \omega_{n-2} \omega_{n+1} |\det C_0|$$

where C_0 is the 4×4 matrix defined by

$$\begin{pmatrix} 1 + \frac{E}{E'} & -1 & 0 & -1 \\ 0 & E' + 2\omega_n & E - E' - 2\omega_n & E' \\ -\omega_{n-1} & \omega_{n-1} + \omega_n - E & -\omega_n & 0 \\ 0 & \frac{E}{E'} & 0 & -1 \end{pmatrix}$$

Finally, by an explicit computation for the determinant of C_0 , we obtain that

$$|\det A_0| = \frac{4E}{E'} (E + E') \omega_{n-2} \omega_{n+1} \left| \omega_n + \frac{E' - E}{4} \right| \leq \rho_0 e^{-L^\beta/4}.$$

Therefore, with the remark that E, E' are positive and $\{\omega_j\}_{j \in \mathbb{Z}}$ are bounded from below by a positive constant, the following condition on ω holds true when L sufficiently large:

$$\left| \omega_n + \frac{E' - E}{4} \right| \leq C e^{-L^\beta/4}. \quad (I)$$

Second case: Now assume that $n-2, n-1 \in \mathcal{Q}_j$ and $n, n+1 \in \mathcal{P}_j$.

Repeating the argument in **First case**, we have a 10×10 system of linear equations

$$A_1 U = b_1$$

where $U = (u(n-2), u(n-1), \dots, u(n+2), v(n-2), v(n-1), \dots, v(n+2))$, $b_1 = (b_1^j)_{1 \leq j \leq 10}$ with $\|b_1\| \leq 10e^{-L^\beta/2}$ and A_1 is the following 10×10 matrix:

$$\begin{pmatrix}
1 & -1 & 0 & 0 & 0 \\
0 & 1 & -1 & 0 & 0 \\
0 & 0 & 1 & -1 & 0 \\
0 & 0 & 0 & 1 & -1 \\
0 & -\omega_{n-1} & \omega_{n-1} + \omega_n - E & -\omega_n & 0 \\
0 & \omega_{n-1} & \omega_n - \omega_{n-1} & -\omega_n & 0 \\
0 & 0 & -\omega_n & \omega_n + \omega_{n+1} - E & -\omega_{n+1} \\
0 & 0 & 0 & \frac{E}{E'} & 0 \\
-\omega_{n-2} & \omega_{n-2} + \omega_{n-1} - E & -\omega_{n-1} & 0 & 0 \\
0 & -\frac{E}{E'} & 0 & 0 & 0
\end{pmatrix}
\begin{pmatrix}
-1 & 1 & 0 & 0 & 0 \\
0 & -1 & 1 & 0 & 0 \\
0 & 0 & 1 & -1 & 0 \\
0 & 0 & 0 & 1 & -1 \\
0 & 0 & 0 & 0 & 0 \\
0 & 0 & -E' & 0 & 0 \\
0 & 0 & 0 & 0 & 0 \\
0 & 0 & 0 & -1 & 0 \\
0 & 0 & 0 & 0 & 0 \\
0 & -1 & 0 & 0 & 0
\end{pmatrix}$$

Denote by $\widetilde{A}_1$ the comatrix of A_1 and put $\rho_1 := \max\{1, \|\widetilde{A}_1\|\}$. As in **First case**, we obtain that

$$|\det A_1| \leq \rho_1 e^{-L^\beta/4}.$$

The determinant of A_1 can be computed as follows:

First, expand this determinant by its sixth and last column and then by its fifth and first column to get

$$|\det A_1| = \omega_{n-2}\omega_{n+1}|\det B_1|$$

where B_1 is the 6×6 matrix defined by

$$\begin{pmatrix}
1 & -1 & 0 & -1 & 1 & 0 \\
0 & 1 & -1 & 0 & -1 & 1 \\
-\omega_{n-1} & \omega_{n-1} + \omega_n - E & -\omega_n & 0 & 0 & 0 \\
\omega_{n-1} & \omega_n - \omega_{n-1} & -\omega_n & 0 & -E' & 0 \\
0 & 0 & \frac{E}{E'} & 0 & 0 & -1 \\
-\frac{E}{E'} & 0 & 0 & -1 & 0 & 0
\end{pmatrix}$$

Second, take the first row of matrix B minus its last row and take the second row minus the fifth row, then expand the determinant of B_1 by its forth and sixth column to obtain

$$|\det A_1| = \omega_{n-2}\omega_{n+1}|\det C_1|$$

where C_1 is the following 4×4 matrix

$$\begin{pmatrix} 1 + \frac{E}{E'} & -1 & 0 & -1 \\ 0 & 1 & -1 - \frac{E}{E'} & 1 \\ -\omega_{n-1} & \omega_{n-1} + \omega_n - E & -\omega_n & -E' \\ \omega_{n-1} & \omega_n - \omega_{n-1} & -\omega_n & -E' \end{pmatrix}$$

Finally, by an explicit computation for the determinant of the matrix C_1 , we infer that

$$|\det A_1| = \frac{4E}{E'} \omega_{n-2} \omega_{n+1} \left| \omega_{n-1} \omega_n - \frac{(E + E')^2}{4} \right| \leq C e^{-L^\beta/4}.$$

Hence, take into account that $\omega_j \geq \alpha_0 > 0$ for all $j \in \mathbb{Z}$ and $E, E' > 0$, we have the following condition on the random variables:

$$\left| \omega_{n-1} \omega_n - \frac{(E + E')^2}{4} \right| \leq C e^{-L^\beta/4} \quad (II)$$

as L sufficiently large.

Third case: Suppose that $n - 2, n - 1 \in \mathcal{Q}_j$, $n \in \mathcal{P}_j$ and $n + 1 \in \mathcal{Q}_{j+1}$. Proceeding as in the previous cases leads us to consider the following system of linear equations

$$A_2 U = b_2$$

where A_2 is the 10×10 matrix defined by

$$\begin{pmatrix} 1 & -1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & -\omega_{n-1} & \omega_{n-1} + \omega_n - E & -\omega_n & 0 \\ 0 & \omega_{n-1} & \omega_n - \omega_{n-1} & -\omega_n & 0 \\ 0 & 0 & -\omega_n & \omega_n + \omega_{n+1} - E & -\omega_{n+1} \\ 0 & 0 & -\omega_n & \omega_n - \omega_{n+1} & \omega_{n+1} \\ -\omega_{n-2} & \omega_{n-2} + \omega_{n-1} - E & -\omega_{n-1} & 0 & 0 \\ 0 & -\frac{E}{E'} & 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} -1 & 1 & 0 & 0 & 0 \\ 0 & -1 & -1 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -E' & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -E' & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 \end{pmatrix}$$

and $|\det A_2| \leq C e^{-L^\beta/4}$.

Then, by an explicit computation, we obtain that

$$|\det A_2| = 4E(E + E')\omega_{n-2}\omega_{n+1}\left|\omega_n + \frac{E' - E}{4}\right| \leq C e^{-L^\beta/4}$$

which immediately yields that

$$\left|\omega_n + \frac{E' - E}{4}\right| \leq C e^{-L^\beta/4} \quad (III)$$

as $L \rightarrow +\infty$.

Forth case: Suppose that $n - 2 \in \mathcal{Q}_j$, $n - 1 \in \mathcal{P}_j$, $n \in \mathcal{Q}_{j+1}$ and $n + 1 \in \mathcal{P}_{j+1}$. Then, $U = (u(n - 2), u(n - 1), \dots, u(n + 2), v(n - 2), v(n - 1), \dots, v(n + 2))$ satisfies the following system of linear equations:

$$A_3 U = b_3$$

where A_3 is the 10×10 matrix defined by

$$\begin{pmatrix} 1 & -1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & 1 & -1 & 0 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & -\omega_{n-1} & \omega_{n-1} + \omega_n - E & -\omega_n & 0 \\ 0 & -\omega_{n-1} & \omega_{n-1} - \omega_n & \omega_n & 0 \\ 0 & 0 & -\omega_n & \omega_n + \omega_{n+1} - E & -\omega_{n+1} \\ 0 & 0 & \omega_n & \omega_{n+1} - \omega_n & -\omega_{n+1} \\ -\omega_{n-2} & \omega_{n-2} + \omega_{n-1} - E & -\omega_{n-1} & 0 & 0 \\ \omega_{n-2} & \omega_{n-1} - \omega_{n-2} & -\omega_{n-1} & 0 & 0 \end{pmatrix} \begin{pmatrix} -1 & 1 & 0 & 0 & 0 \\ 0 & 1 & -1 & 0 & 0 \\ 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & 1 & -1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & -E' & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -E' & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & -E' & 0 & 0 & 0 \end{pmatrix}$$

and $|\det A_3| \leq C e^{-L^\beta/4}$.

We compute

$$|\det A_3| = E E' \omega_{n-2} \times \omega_{n+1} \times \left|4\omega_{n-1} + E' - E\right| \times \left|E' - E + 4\omega_n\right|.$$

Hence, at least one of the two following conditions on ω must be satisfied:

$$\bullet \left|\omega_{n-1} + \frac{E' - E}{4}\right| \leq C e^{-L^\beta/8}, \quad (IV)$$

$$\bullet \left| \omega_n + \frac{E' - E}{4} \right| \leq C e^{-L^\beta/8}.$$

Fifth case: Assume that $n - 2, n - 1, n, n + 1 \in \mathcal{P}_j$. Hence, by **Step 1**, $n + 2 \in \mathcal{Q}_j$. We consider four points $n - 1, n, n + 1, n + 2$ and come back to **First case**.

Last case: Suppose that $n - 2 \in \mathcal{Q}_j$, $n - 1, n \in \mathcal{P}_j$, and $n + 1 \in \mathcal{Q}_{j+1}$. We consider the point $n + 2$.

If $n + 2$ belongs to $\mathcal{Q}_{j+1}$, we consider four points $n - 1, n, n + 1, n + 2$ and come back to **Second case** (by the equivalent role of $\mathcal{P}$ and $\mathcal{Q}$). Otherwise $n + 2$ belongs to $\mathcal{P}_{j+1}$. In this case, we consider four points $n - 1, n, n + 1, n + 2$ and come back to **Third case**.

To conclude, the random variables ω_j must satisfy at least cL^β conditions of one of the types (I)-(IV). From the fact that ω_n are i.i.d. and possess a bounded density, the conditions (I)-(IV) imply that the event (*) can occur for a given partition $\mathcal{P}$ and $\mathcal{Q}$ with a probability at most, $(e^{-cL^\beta})^{L^\beta} = e^{-cL^{2\beta}}$, for some $c > 0$. Hence,

$$\mathbb{P}^* \leq 2^L e^{-cL^{2\beta}} \leq e^{-\tilde{c}L^{2\beta}}$$

as the number of partitions is bounded by 2^L and $\beta > 1/2$.

We therefore have Lemma 3.3 proved. $\square$

Remark: Thanks to the equality (3.3), it is not hard to derive the following estimate for the model (1.1),

$$\frac{\Delta E}{2\beta_0} |\Lambda|^{-1/2} \leq \|\nabla_\omega(E(\omega) - E'(\omega))\|_2 \leq \|\nabla_\omega(E(\omega) - E'(\omega))\|_1 \quad (3.21)$$

provided that $|E(\omega) - E| + |E'(\omega) - E'| \leq e^{-L^\beta}$ and ΔE is the distance between E and E' .

The above estimate reads that the l^1 -distance of the gradients of $E(\omega)$ and $E'(\omega)$ is bounded from below by a positive term that is not exponentially small w.r.t. the length of the interval Λ .

Now, let $c_1 = c_2$ in Lemma 3.3.

Under the hypotheses in Lemma 3.3, the estimate (3.21) and Lemma 3.5 immediately imply that

$$\frac{\Delta E}{2\beta_0} |\Lambda|^{-1/2} \leq C |\Lambda| e^{-|\Lambda|^\beta}$$

that is impossible if $|\Lambda|$ is sufficiently large. Hence, for $c_1 = c_2$, $\mathbb{P}^*$ is equal to 0.

Finally, we would like to note that an estimate as (3.21) for the discrete Anderson model only holds true for two distinct energies sufficiently far apart from each other. Moreover, that kind of estimate enable us to prove the decorrelation estimate for the discrete Anderson model in *any* dimension (c.f. Lemma 2.4 in [9]). But it is not the case for the model (1.1).

4 Comment on the lower bound of the r.v.'s

In this section, we want to discuss how to relax the hypothesis on the lower bound of random variables $\{\omega_j\}_{j \in \mathbb{Z}}$.

Assume that all r.v.'s ω_j are only non-negative instead of being bounded from below by a positive constant.

Thanks to the Taylor expansion, the common distribution function $F(t)$ of $\{\omega_j\}_{j \in \mathbb{Z}}$ always satisfies the following equality:

$$F(t) = \rho(0)t + o(t) \quad (4.1)$$

near 0 where ρ is the density function of the r.v.'s $\{\omega_j\}_{j \in \mathbb{Z}}$.

Hence,

$$F(t) \geq \frac{\rho(0)}{2}t \quad (4.2)$$

near 0 provided that $\rho(0)$ is positive.

We will show that decorrelation estimates can be obtained by using (4.2) i.e., by using the assumption $\rho(0) > 0$.

Pick $\Lambda = [-L, L]$ as usual and β a fixed number in $(0, 1)$ which will be chosen later. Then, we have $2L + 1 - L^\beta$ sub-intervals of length L^β in Λ .

Using the independence of the r.v.'s ω_j , we infer that

$$\begin{aligned} \mathbb{P}^1 &:= \mathbb{P}(\exists \text{ an interval } J \text{ of length } L^\beta \text{ in } \Lambda \text{ such that } \omega_j \geq e^{-(\log L)^\delta} \forall j \in J) \\ &\leq (2L + 1 - L^\beta)(1 - \mathbb{P}(\omega_0 < e^{-(\log L)^\delta}))^{L^\beta} \leq 2L(1 - F(e^{-(\log L)^\delta}))^{L^\beta} \end{aligned}$$

where δ is an arbitrary, fixed number in $(0, 1/3)$.

Next, we put $x := -F(e^{-(\log L)^\delta}) \in (-1, 0)$, and rewrite the quantity $(1 - F(e^{-(\log L)^\delta}))^{L^\beta}$ in the form $((1 + x)^{1/x})^{xL^\beta}$.

We observe that the function $f(x) = (1 + x)^{1/x}$ is decrease in $(-1, 0)$ and $\lim_{x \rightarrow 0} f(x) = e$, hence $f(x) \geq e \forall x \in (-1, 0)$.

Combining this with (4.2), for L large enough, we get

$$\mathbb{P}^1 \leq 2Le^{-L^\beta F(e^{-(\log L)^\delta})} \leq 2Le^{-c_0 L^\beta e^{-(\log L)^\delta}} \lesssim 2Le^{-cL^{\beta_1}} \lesssim e^{-cL^{\beta_1}} \quad (4.3)$$

where c_0, c, β_1 are positive constants and β_1 is smaller than β .

In other words, with a probability $1 - e^{-cL^{\beta_1}}$, there exists a sub-interval J of side length L^β of Λ such that

$$\omega_j \geq e^{-(\log L)^\delta} \quad \forall j \in J.$$

We can therefore use the same argument in the proof of Lemma 3.2 to get:

Lemma 4.1. *Let $\Lambda = [-L, L]$ be a large cube in $\mathbb{Z}$ and ϵ be a fixed number in the interval $(0, \beta)$.*

Then, with the probability $1 - e^{-cL^{\beta_1}}$, there exists a point k_0 in Λ and a positive constant c_ϵ such that $u^2(k) + u^2(k+1) \geq e^{-L^\beta}$ for all $|k - k_0| \leq c_\epsilon L^{\beta(\beta-\epsilon)}$ as L is large enough.

Hence, the upper bound for the probability $\mathbb{P}$ in Lemma 3.3 is now $B := c2^L e^{-cL^{\beta^2+\beta/2}}$ if we choose ϵ to be $\beta/2$ in Lemma 4.1.

Observe that $\beta^2 + \beta/2 > 1$ provided that $1 > \beta > \frac{\sqrt{17}-1}{4}$. With this choice of β , the upper bound B is exponentially small as L tends to plus infinity. Then we can proceed again as in Lemma 3.3 and remark that the error term $e^{-cL^{\beta_1}} \ll (l/L)^2 e^{(\log L)^\beta}$, we obtain again a decorrelation estimate for the model (1.1):

Theorem 4.1. *For $\beta \in \left(\frac{\sqrt{17}-1}{4}, 1\right)$, $\alpha \in (0, 1)$ and $l = O(L^\alpha)$, we have*

$$\mathbb{P}\left(\left\{\begin{array}{l} \sigma(H_\omega(\Lambda_l)) \cap (E + L^{-1}(-1, 1)) \neq \emptyset \\ \sigma(H_\omega(\Lambda_l)) \cap (E' + L^{-1}(-1, 1)) \neq \emptyset \end{array}\right\}\right) \leq C(l/L)^2 e^{(\log L)^\beta} = o(l/L).$$

when L is sufficiently large.

5 More than two distinct energies

In this section, we would like to show that, following an argument in [9], we can use Theorem 1.2 to prove the asymptotic independence for any fixed number of point processes, i.e., for a fixed number $n \geq 2$ and a finite sequence of fixed energies $\{E_i\}_{1 \leq i \leq n}$ in the localized regime, n point processes $\Xi(\xi, E_i, \omega, \Lambda)$ defined as in (1.3) converge to n independent Poisson processes. For example, we will prove in detail the case of $n = 3$ with three distinct energies E, E', E'' .

Consider non-empty compact intervals $(U_j)_{1 \leq j \leq J}$, $(U'_j)_{1 \leq j \leq J'}$, $(U''_j)_{1 \leq j \leq J''}$ and integers $(k_j)_{1 \leq j \leq J}$, $(k'_j)_{1 \leq j \leq J'}$, $(k''_j)_{1 \leq j \leq J''}$ as in Theorem 1.3.

Using notations in [9], we pick L and l such that $(2L+1) = (2l+1)(2l'+1)$, $cL^\alpha \leq l \leq L^\alpha/c$ for some $\alpha \in (0, 1)$ and $c > 0$.

We decompose the interval $\Lambda := [-L, L] = \bigcup_{|\gamma| \leq l'} \Lambda_l(\gamma)$ where $\Lambda_l(\gamma) := (2l+1)\gamma + \Lambda_l$.

Next, for $\Lambda' \subset \Lambda$ and $U \subset \mathbb{R}$, consider the r.v.'s

$$X(E, U, \Lambda') := \begin{cases} 1 & \text{if } H_\omega(\Lambda') \text{ has at least one eigenvalue} \\ & \text{in } E + (\nu(E)|\Lambda'|)^{-1}U, \\ 0 & \text{otherwise} \end{cases} \quad (5.1)$$

and put $\Sigma(E, U, l) := \sum_{|\gamma| \leq l'} X(E, U, \Lambda_l(\gamma))$.

First of all, [Lemma 3.2, [9]] is the first ingredient of the proof which tell us that we can actually reduce our problem to consider eigenvalues associated to smaller intervals. This lemma is still true in the three -energy-case.

Hence, to complete the proof of the stochastic independence w.r.t. three processes, the only thing left to do is show that the joint finite dimensional

distribution function of three point processes tends to the product of three finite dimensional distributions associated to each of three energies, i.e.,

$$\mathbb{P} \left(\left\{ \begin{array}{l} \omega; \quad \Sigma(E, U_1, l) = k_1, \quad \dots, \Sigma(E, u(j), l) = k_J \\ \Sigma(E', U'_1, l) = k'_1, \quad \dots, \Sigma(E', U'_{j'}, l) = k'_{j'} \\ \Sigma(E'', U''_1, l) = k''_1, \quad \dots, \Sigma(E'', U''_{j''}, l) = k''_{j''} \end{array} \right\} \right) \quad (5.2)$$

should be well approximated by the product

$$\begin{aligned} & \mathbb{P} \left(\left\{ \begin{array}{l} \omega; \quad \Sigma(E, U_1, l) = k_1 \\ \vdots \\ \Sigma(E, u(j), l) = k_J \end{array} \right\} \right) \times \mathbb{P} \left(\left\{ \begin{array}{l} \omega; \quad \Sigma(E', U'_1, l) = k'_1 \\ \vdots \\ \Sigma(E', U'_{j'}, l) = k'_{j'} \end{array} \right\} \right) \\ & \times \mathbb{P} \left(\left\{ \begin{array}{l} \omega; \quad \Sigma(E'', U''_1, l) = k''_1 \\ \vdots \\ \Sigma(E'', U''_{j''}, l) = k''_{j''} \end{array} \right\} \right) \end{aligned}$$

as L goes to infinity.

By a standard criterion of the convergence of point processes (c.f. e.g. Theorem 11.1.VIII, [3]), the above statement holds true if the following quantity vanishes for all real numbers $t_j, t'_{j'}, t''_{j''}$:

$$\begin{aligned} & \mathbb{E} \left(e^{-\sum_{j=1}^J t_j \Sigma(E, u(j), l) - \sum_{j'=1}^{J'} t'_{j'} \Sigma(E', U'_{j'}, l) - \sum_{j''=1}^{J''} t''_{j''} \Sigma(E'', U''_{j''}, l)} \right) - \\ & \mathbb{E} \left(e^{-\sum_{j=1}^J t_j \Sigma(E, u(j), l)} \right) \mathbb{E} \left(e^{-\sum_{j'=1}^{J'} t'_{j'} \Sigma(E', U'_{j'}, l)} \right) \mathbb{E} \left(e^{-\sum_{j''=1}^{J''} t''_{j''} \Sigma(E'', U''_{j''}, l)} \right) \end{aligned}$$

as L goes to infinity.

Remark that, from the fact that $\{\Lambda(\gamma)\}_{|\gamma| \leq l'}$ are pairwise disjoint intervals, operators $\{H_\omega(\Lambda(\gamma))\}_{|\gamma| \leq l'}$ are independent operator-valued r.v.'s.

We thus have:

$$\begin{aligned} & \mathbb{E} \left(e^{-\sum_{j=1}^J t_j \Sigma(E, u(j), l) - \sum_{j'=1}^{J'} t'_{j'} \Sigma(E', U'_{j'}, l) - \sum_{j''=1}^{J''} t''_{j''} \Sigma(E'', U''_{j''}, l)} \right) \\ & = \mathbb{E} \prod_{|\gamma| \leq l'} \left(e^{-\sum_j t_j X(E, u(j), \gamma) - \sum_{j'} t'_{j'} X(E', U'_{j'}, \gamma) - \sum_{j''} t''_{j''} X(E'', U''_{j''}, \gamma)} \right) \\ & = \prod_{|\gamma| \leq l'} \mathbb{E} \left(e^{-\sum_j t_j X(E, u(j), \gamma) - \sum_{j'} t'_{j'} X(E', U'_{j'}, \gamma) - \sum_{j''} t''_{j''} X(E'', U''_{j''}, \gamma)} \right). \end{aligned}$$

Our goal is to approximate terms of the following form

$$\mathbb{E} \left(e^{-\sum_j t_j X(E, u(j), \gamma) - \sum_{j'} t'_{j'} X(E', U'_{j'}, \gamma) - \sum_{j''} t''_{j''} X(E'', U''_{j''}, \gamma)} \right)$$

by the product

$$\prod_j^J \mathbb{E} e^{-t_j X(E, u(j), \gamma)} \prod_{j'}^{J'} \mathbb{E} e^{-t_{j'} X(E', U'_{j'}, \gamma)} \prod_{j''}^{J''} \mathbb{E} e^{-t_{j''} X(E'', U''_{j''), \gamma)}$$

as L large enough with the remark that these r.v.'s are not independent. To illustrate our computation for that, we will consider just three arbitrary Bernoulli r.v.'s X_1, X_2, X_3 and we will compute explicitly $A := \mathbb{E}(e^{\sum_{i=1}^3 a_i X_i})$.

$$\begin{aligned} A &= \mathbb{P}(X_1 = X_2 = X_3 = 0) + \sum_{i=1}^3 e^{a_i} \mathbb{P} \left(\begin{array}{l} X_i = 1, \\ X_j = 0 \ \forall j \neq i \end{array} \right) \\ &+ \sum_{i < j} e^{a_i + a_j} \mathbb{P} \left(\begin{array}{l} X_i = X_j = 1, \\ X_k = 0; k \neq i, j \end{array} \right) + e^{a_1 + a_2 + a_3} \mathbb{P} \left(\bigcap_{i=1}^3 X_i \right). \end{aligned}$$

First of all, we rewrite

$$\begin{aligned} \mathbb{P}(X_1 = X_2 = X_3 = 0) &= 1 - \sum \left(\begin{array}{l} X_i = 1, \\ X_j = 0 \ \forall j \neq i \end{array} \right) \\ &- \sum_{i < j} \mathbb{P} \left(\begin{array}{l} X_i = X_j = 1, \\ X_k = 0; k \neq i, j \end{array} \right) - \mathbb{P} \left(\bigcap_{i=1}^3 X_i \right) \end{aligned}$$

and obtain that

$$\begin{aligned} A &= 1 + \sum (e^{a_i} - 1) \left(\begin{array}{l} X_i = 1, \\ X_j = 0 \ \forall j \neq i \end{array} \right) \\ &+ \sum_{i < j} (e^{a_i + a_j} - 1) \mathbb{P} \left(\begin{array}{l} X_i = X_j = 1, \\ X_k = 0; k \neq i, j \end{array} \right) + (e^{a_1 + a_2 + a_3} - 1) \mathbb{P} \left(\bigcap_{i=1}^3 X_i \right). \end{aligned}$$

Next, use

$$\begin{aligned} \mathbb{P} \left(\begin{array}{l} X_i = 1, \\ X_j = 0 \ \forall j \neq i \end{array} \right) &= \mathbb{P}(X_i = 1) - \mathbb{P} \left(\begin{array}{l} X_i = X_{i+1} = 1, \\ X_{i+2} = 0 \end{array} \right) \\ &- \mathbb{P} \left(\begin{array}{l} X_i = X_{i-1} = 1, \\ X_{i+1} = 0 \end{array} \right) - \mathbb{P} \left(\bigcap_{i=1}^3 X_i \right) \end{aligned}$$

to get

$$\begin{aligned} A &= 1 + \sum (e^{a_i} - 1) \mathbb{P}(X_i = 1) + \sum_{i < j} (e^{a_i} - 1)(e^{a_j} - 1) \mathbb{P} \left(\begin{array}{l} X_i = X_j = 1, \\ X_k = 0 \ k \neq i, j \end{array} \right) \\ &+ \left(e^{\sum_{i=1}^3 a_i} - 1 - \sum_{i=1}^3 (e^{a_i} - 1) \right) \mathbb{P} \left(\bigcap_{i=1}^3 X_i \right). \end{aligned}$$

Using a similar expansion for all terms of the form $\mathbb{P}\left(\begin{smallmatrix} X_i = X_j = 1, \\ X_k = 0 \ k \neq i, j \end{smallmatrix}\right)$, we obtain the following expression of $\mathbb{E}\left(e^{\sum_{i=1}^3 a_i X_i}\right)$

$$\begin{aligned} A &= 1 + \sum (e^{a_i} - 1) \mathbb{P}(X_i = 1) + \sum_{i < j} (e^{a_i} - 1)(e^{a_j} - 1) \mathbb{P}(X_i = X_j = 1) \\ &\quad + \prod_{i=1}^3 (e^{a_i} - 1) \mathbb{P}\left(\bigcap_{i=1}^3 X_i\right). \end{aligned}$$

On the other hand, from the observation that

$$\mathbb{E}e^{a_j X_j} = 1 + (e^{a_j} - 1) \mathbb{P}(X_j = 1), \quad \forall j = \{1, 2, 3\},$$

one multiplies the three equalities above to get

$$\begin{aligned} \prod_{i=1}^3 \mathbb{E}(e^{a_i X_i}) &= 1 + \sum (e^{a_i} - 1) \mathbb{P}(X_i = 1) \\ &\quad + \sum_{i < j} (e^{a_i} - 1)(e^{a_j} - 1) \mathbb{P}(X_i = 1) \mathbb{P}(X_j = 1) + \prod_{i=1}^3 (e^{a_i} - 1) \prod_{i=1}^3 \mathbb{P}(X_i = 1). \end{aligned}$$

Hence, the difference between the expectation of $e^{\sum_{i=1}^3 a_i X_i}$ and the product of expectations $e^{a_i X_i}$ is:

$$\begin{aligned} \mathbb{E}\left(e^{\sum_{i=1}^3 a_i X_i}\right) - \prod_{i=1}^3 \mathbb{E}(e^{a_i X_i}) &= \\ &= \sum_{i < j} (e^{a_i} - 1)(e^{a_j} - 1) [\mathbb{P}(X_i = X_j = 1) - \mathbb{P}(X_i = 1) \mathbb{P}(X_j = 1)] \\ &\quad + \prod_{i=1}^3 (e^{a_i} - 1) \left[\mathbb{P}\left(\bigcap_{i=1}^3 X_i\right) - \prod_{i=1}^3 \mathbb{P}(X_i = 1) \right]. \end{aligned}$$

Thanks to the decorrelation estimate and the Wegner estimate, we obtain

$$\begin{aligned} &\mathbb{P}(X_i = X_j = 1) + \mathbb{P}(X_i = 1) \mathbb{P}(X_j = 1) \\ &\leq C(l/L)^2 (e^{(\log L)^\beta} + 1) \leq C(l/L)^{1+\rho} \end{aligned}$$

and

$$\begin{aligned} &\mathbb{P}\left(\bigcap_{i=1}^3 X_i\right) + \mathbb{P}(X_1 = 1) \mathbb{P}(X_2 = 1) \mathbb{P}(X_3 = 1) \\ &\leq C(l/L)^2 (e^{(\log L)^\beta} + (l/L)) \leq C(l/L)^{1+\rho} \end{aligned}$$

where $\rho \in (0, 1)$.

To sum up, we get that:

$$\mathbb{E}(e^{\sum_{i=1}^3 a_i X_i}) = \prod_{i=1}^3 \mathbb{E}(e^{a_i X_i}) + O((l/L)^{1+\rho}).$$

In other words, $\mathbb{E}(e^{\sum_{i=1}^3 a_i X_i})$ is well approximated by $\prod_{i=1}^3 \mathbb{E}(e^{a_i X_i})$ with a small error term of the magnitude $(l/L)^{1+\rho}$. It is easy to see that we can obtain the same result for an arbitrarily finite number of Bernoulli r.v.'s. Hence, for each $|\gamma| \leq l'$, we have:

$$\begin{aligned} & \mathbb{E} \left(e^{-\sum_j t_j X(E, u(j), \gamma) - \sum_{j'} t'_{j'} X(E', U'_{j'}, \gamma) - \sum_{j''} t''_{j''} X(E'', U''_{j''), \gamma)} \right) \\ &= \prod_j \mathbb{E} e^{-t_j X(E, u(j), \gamma)} \prod_{j'} \mathbb{E} e^{-t'_{j'} X(E', U'_{j'}, \gamma)} \prod_{j''} \mathbb{E} e^{-t''_{j''} X(E'', U''_{j''), \gamma)} \\ &\times (1 + O(l/L)^{1+\rho}). \end{aligned}$$

Besides, we have similar formulas for the expectation of $e^{-\sum_j t_j X(E, u(j), \gamma)}$, $e^{-\sum_j t_j X(E', U'_{j'}, \gamma)}$ and $e^{-\sum_j t_j X(E'', U''_{j''), \gamma)}$.

We therefore infer that:

$$\begin{aligned} & \mathbb{E} \left(e^{-\sum_j t_j X(E, u(j), \gamma) - \sum_{j'} t'_{j'} X(E', U'_{j'}, \gamma) - \sum_{j''} t''_{j''} X(E'', U''_{j''), \gamma)} \right) \\ &= \mathbb{E} e^{-\sum_j t_j X(E, u(j), \gamma)} \mathbb{E} e^{-\sum_{j'} t'_{j'} X(E', U'_{j'}, \gamma)} \mathbb{E} e^{-\sum_{j''} t''_{j''} X(E'', U''_{j''), \gamma)} \\ &\times (1 + O(l/L)^{1+\rho}). \end{aligned}$$

An important observation is that we only have $l' = O(L/l)$ sub-intervals of the side length l of the interval Λ . Hence, as multiplying all above equalities side by side over γ , we obtain that

$$\mathbb{E} \left(e^{-\sum_{j=1}^J t_j \Sigma(E, u(j), l) - \sum_{j'=1}^{J'} t'_{j'} \Sigma(E', U'_{j'}, l) - \sum_{j''=1}^{J''} t''_{j''} \Sigma(E'', U''_{j''), l)} \right)$$

is equal to the product of

$$\mathbb{E} \left(e^{-\sum_{j=1}^J t_j \Sigma(E, u(j), l)} \right) \mathbb{E} \left(e^{-\sum_{j'=1}^{J'} t'_{j'} \Sigma(E', U'_{j'}, l)} \right) \mathbb{E} \left(e^{-\sum_{j''=1}^{J''} t''_{j''} \Sigma(E'', U''_{j''), l)} \right)$$

and an error term of the form

$$(1 + x^{1+\rho})^{1/x}$$

which tends to 1 as L goes to infinity with $x = O(l/L)$.

Hence, the stochastic independence for three point processes w.r.t. three distinct energies is proved.

Acknowledgements. This work would have been impossible without great help of my advisor, Prof. Frédéric Klopp. The author gratefully acknowledges the collaboration and various discussions with him.

References

- [1] Alexander Figotin and Abel Klein, *Localization of electromagnetic and acoustic waves in random media. Lattice models*, J. Statist. Phys., 76(3-4):9851003, 1994.
- [2] Michael Aizenman, Jeffrey H. Schenker, Roland M. Friedrich, and Dirk Hundertmark, *Finite-volume fractional-moment criteria for Anderson localization*, Comm. Math. Phys., 224(1):219-253, 2001. Dedicated to Joel L. Lebowitz.
- [3] D.J. Daley, D. Vere-Jones, *An Introduction to the Theory of Point Processes. Vol. II. General Theory and Structure*, 2nd edition, Probability and its applications (New York). Springer, New York (2008)
- [4] Peter D. Hislop, *Lectures on random Schrödinger operators*, In Fourth Summer School in Analysis and Mathematical Physics, volume 476 of Contemp. Math., pages 411-31. Amer. Math. Soc., Providence, RI, 2008.
- [5] Werner Kirsch and Bernd Metzger, *The integrated density of states for random Schrödinger operators*, In Spectral theory and mathematical physics: a Festschrift in honor of Barry Simons 60th birthday, volume 76 of Proc. Sympos. Pure Math., pages 649-696. Amer. Math. Soc., Providence, RI, 2007.
- [6] Dong Miao, *Eigenvalue statistics for lattice Hamiltonian of off-diagonal disorder*, J. Stat. Phys (2011), 143: 509-522 DOI 10.1007/s10955-011-0190-2.
- [7] Nariyuki Minami, *Local fluctuation of the spectrum of a multidimensional Anderson tight binding model*, Comm. Math. Phys. Vol. 177, 709-725 (1996).
- [8] François Germinet and Frédéric Klopp, *Spectral statistics for random Schrödinger operators in the localized regime*, To appear in Jour. Eur. Math. Soc.
- [9] Frédéric Klopp, *Decorrelation estimates for the eigenvalues of the discrete Anderson model in the localized regime*, Comm. Math. Phys. Vol. 303, pp. 233-260 (2011).
- [10] Tosio Kato, *Perturbation Theory for Linear Operators*. Springer-Verlag, Berlin, 1995. Reprint of 1980 edition.
- [11] Ivan Veselić, *Existence and regularity properties of the integrated density of states of random Schrödinger operators*, volume 1917 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, 2008.

(Trinh Tuan Phong) Laboratoire Analyse, Géométrie et Applications,
UMR 7539, Institut Galilée, Université Paris 13,
99 avenue J.B. Clément, 93430 Villetaneuse, France
Email: trinh@math.univ-paris13.fr